



Gambaruto, A., & van Loon, R. (2018). Three-Dimensional Simulations of Idealised Lymphangion. In *BioMedEng18 Proceedings: The UK's largest gathering of Biomedical Engineers Medical Engineers and Bioengineers - 6-7 September 2018, Imperial College London* (pp. 386). BioMedEng.

Peer reviewed version

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the final published version of the article (version of record). It first appeared online via BioMedEng at [https://research-information.bris.ac.uk/pure/files/161872649/BioMedEng\\_gambaruto.pdf](https://research-information.bris.ac.uk/pure/files/161872649/BioMedEng_gambaruto.pdf) . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

# BioMedEng18

## Three-dimensional Simulations of Idealised Lymphangion

**A.M. Gambaruto<sup>1</sup>, R. van Loon<sup>2</sup>**

<sup>1</sup> Queen's Building, University of Bristol, BS8 1TR, Bristol. <sup>2</sup> Bay Campus, Swansea University, SA1 8EN, Swansea.

### Introduction

The lymphatic system is responsible for collecting fluid leaked from blood vessels, returning it to the blood system to maintain overall fluid balance. Lymphatic vessels also fulfill important immune functions by transporting leukocytes and antigens [1,2,3]. The functional units of collecting lymphatic vessels are called the lymphangions [2,4]. Some experimental and computational studies exist, however much still needs learnt about the transport mechanisms.

### Methods

An idealised geometry of a lymphangion unit is considered. This consists of a deforming cylinder with valves at either end. The flow is considered to be periodic. The Moving Particle Semi-implicit (MPS) method is employed, since this has been shown to be effective in simulation of micro-circulation [5]. Periodic boundary conditions are employed in order to investigate time integrated measures.

### Results & Discussion

A parametric study is carried out, varying geometry of the lymphangion and the pulsatility of the lymph. We study the redistribution of particles as they traverse the valve constriction, looking at the forces imposed on cells and the transport properties.

### References

1. Moore Jr JE, Bertram CD. Lymphatic System Flows. *Annual Review of Fluid Mechanics*. 2018; 50:459-482.
2. Margaris KN, Black RA. Modelling the lymphatic system: challenges and opportunities. *Journal of the Royal Society Interface*. 2012; 9(69):601-612.
3. Bazigou E, Wilson JT, Moore Jr JE. Primary and secondary lymphatic valve development: molecular, functional and mechanical insights. *Microvascular research*. 2014; 96:38-45.
4. Russo E, Nitschké M, Halin C. Dendritic cell interactions with lymphatic endothelium. *Lymphatic research and biology*. 2013; 11(3):172-182.
5. Gambaruto AM. Computational haemodynamics of small vessels using the moving particle semi-implicit (MPS) method. *Journal of Computational Physics*. 2015;302:68-96.

### Acknowledgments

The authors declare no conflicts of interests.